



## Native/Wildflower Seed Mixes

### Meadow Mixture

Provides a colourful flower display along with soil stabilization, and wildlife habitat for open areas.

Catchfly 8%  
 New England Aster 3%  
 Canada Milk Vetch 3%  
 Lance Leaved Coreopsis 10%  
 Dense Blazingstar 5%  
 Bergamot 2%  
 Black Eyed Susan 8%  
 Riddell's Goldenrod 3%  
 Ohio Spiderwort 3%  
 Blue Vervain 8%  
 Boneset 2%  
 Big Bluestem 12%  
 Little Bluestem 10%  
 Canada Wild Rye 13%  
 Indian Grass 10%

Seeding rate - 21 - 23 lbs/acre, 22 - kg/hectare  
 1/2 lb/1000 sq.ft., 250 gm/90 sq.m

### Wet Meadow Mixture

Contains a blend of species to provide colour, diversity of plant growth and potential wildlife habitat for damp, open locations.

Canada Bluejoint 3%  
 Fowl Bluegrass 18%  
 Fowl Mannagrass 10%  
 Soft Rush 5%  
 New England Aster 3%  
 Wool Grass 5%  
 Black Eyed Susan 4%  
 Boneset 2%  
 Blue Vervain 5%  
 Big Blue Stem 12%  
 Virginia Wild Rye 18%  
 Riverbank Wild Rye 15%

Seeding rate - 21 - 23 lbs/acre, 22 - 25 kg/hectare  
 1/2 lb/1000 sq.ft., 250 gm/ 90 sq.m

### Perennial Mixture

Offers a colourful display of perennial plants.

Black Eyed Susan 10%  
 Coreopsis Lance Leaved 10%  
 Catchfly 15%  
 Purple Coneflower 5%  
 Biannual Evening Primrose 5%  
 Wild Blue Lupine 10%  
 Shasta Daisy 15%  
 Sweet William Double 15%  
 Siberian Wallflower 15%

Seeding rate - 7 - 9 lbs/acre, 8 - 10 kg/hectare  
 1/4 lb/1000 sq.ft., 100 gm/90 sq.m

### Wetland Mixture

Offers a naturalization blend suited to predominantly wet soil conditions.

Big Bluestem 15%  
 Canada Bluejoint 8%  
 Fox Sedge 8%  
 Hard Stemmed Bulrush 5%  
 American Mannagrass 9%  
 Green Bulrush 5%  
 Soft Rush 5%  
 Lurid (Shallow) Sedge 5%  
 Boneset 3%  
 Blue Vervain 8%  
 Virginia Wild Rye 19%  
 Riverbank Wild Rye 10%

Seeding rate - 21 - 23 lbs/acre, 22 - 25 kg/hectare  
 1/2 lb/1000 sq. ft., 250 gm/90 sq.m

### Pond Mixture

Provides soil stabilization and potential wildlife habitat for areas surrounding swamps, bogs and pond locations.

Giant Bur Reed 18%  
 Arrow Arum 20%  
 Soft Stem Bulrush 9%  
 Blue Flag 5%  
 Wool Grass 10%  
 Buttonbush 10%  
 Cosmos (Bristly) Sedge 5%  
 Lurid (Shallow) Sedge 10%  
 Purple Stemmed Aster 5%  
 Blue Vervain 8%

Seeding rate - 21 - 23 lbs/acre, 22 - 25 kg/hectare  
 1/2 lb/1000 sq. ft., 250 gm/90 sq.m

### Wildflower Butterfly Mixture

Offers a cross-section of wildflowers whose nectar and showy display will attract butterflies.

Bachelor Buttons/Cornflower 10%  
 Butterfly Milkweed 3%  
 Catchfly 10%  
 Lemon Mint 10%  
 Grey Headed Coneflower 5%  
 Coneflower(Purple) 5%  
 Coreopsis (Plains) 10%  
 Coreopsis Lance Leaved 10%  
 Cosmos 10%  
 Wild Bergamot 2%  
 New England Aster 5%  
 Black Eyed Susan 10%  
 Wild Blue Lupine 10%

Seeding rate - 21 - 23 lbs/acre, 22 - 25 kg/hectare  
 1/2 lb/1000 sq. ft., 250 gm/90 sq.m

### Trail Mixture

Provides a blend of grasses and wildflowers for soil stabilization and wildlife cover adapted to drier soil conditions.

Sand Dropseed 18%  
 Little Bluestem 18%  
 Canada Wild Rye 20%  
 Indian Grass 17%  
 Black Eyed Susan 10%  
 Bergamot 3%  
 Wild Blue Lupine 3%  
 Purple Coneflower 3%  
 Showy Evening Primrose 3%  
 Showy Tick Trefoil 5%

Seeding rate - 21 - 23 lbs/acre, 22 - 25 kg/hectare  
 1/2 lb/1000 sq. ft., 250 gm/90 sq.m

### Prairie Mixture (Medium Height)

Offers a blend of grass species providing soil stabilization and wildlife habitat for flat open areas.

Little Bluestem 10%  
 Canada Wild Rye 10%  
 Indian Grass 20%  
 Side Oats Grama 15%  
 Big Bluestem 20%  
 Switch Grass 15%  
 Prairie Dropseed 5%  
 Prairie Cordgrass 5%

Seeding rate - 21 - 23 lbs/acre, 22 - 25 kg/hectare  
 1/2 lb/1000 sq. ft., 250 gm/90 sq.m



### USEFUL CALCULATIONS

1 Acre = 43,560 Square Feet  
 1 Acre = 4047.86 Square Metres  
 1 KG = 2.2046 lbs  
 1 Square Foot = .09 Square Metres

Mixes should be seeded at the recommended rates. Soil should be worked to ensure good seed-to-soil contact. Annual ryegrass may be used as a nurse crop or for erosion control at the rate of 20 lbs/acre (22 kg/ha). The ideal planting time is in the fall, mid-October to late November. Early spring sowing is less effective. Environmental conditions, such as moisture level, will impact seed germination and in some cases delay establishment until the following year. Rodent and bird damage may be a possibility and will impact seed establishment. Seed mixtures may vary slightly from those listed depending on ingredient availability. We reserve the right to substitute varieties of equal ecological value as required. The buyer acknowledges that acceptance of the seed herein and that the limitations described above are conditions of sale and constitute the entire agreement between the parties regarding warranty and liability.